

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

A02

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/87, 15/86, 15/67, A61K 48/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 97/39135</b>
			(43) International Publication Date: 23 October 1997 (23.10.97)
<p>(21) International Application Number: PCT/US97/05325</p> <p>(22) International Filing Date: 1 April 1997 (01.04.97)</p> <p>(30) Priority Data: 60/015,790 17 April 1996 (17.04.96) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 60/015,790 (CIP) Filed on 17 April 1996 (17.04.96)</p> <p>(71) Applicant (for all designated States except US): BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West 7th Street, Austin, TX 78701 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): CRISTIANO, Richard, J. [US/US]; 2718 Aspen Lane, Pearland, TX 77584 (US). NGUYEN, Dao [CA/CA]; 120 Thornton, Town of Mont Royal, Québec H3P 1H6 (CA).</p> <p>(74) Agent: HIGHLANDER, Steven, L.; Arnold, White &amp; Durkee, P.O. Box 4433, Houston, TX 77210 (US).</p>			<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
(54) Title: ENHANCED EXPRESSION OF TRANSGENES			
(57) Abstract			
<p>The present invention provides for improved methods of gene transfer, both <i>in vitro</i> and <i>in vivo</i>. By treating neoplastic cells with a DNA-damaging agent prior to transduction with a transgene, the expression of the transgene is improved up to about 3-fold over the expression seen in the absence of the DNA-damaging agent treatment. This effect is not dependent on the tumor cell type, the method of DNA transduction or type of DNA-damaging agent. The effect is most dramatic when the transduction is performed about 1-3 days following treatment with the DNA-damaging agent.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## DESCRIPTION

### ENHANCED EXPRESSION OF TRANSGENES

5

### BACKGROUND OF THE INVENTION

The government may own rights in the present invention pursuant to funding of research under NIH Grant No. CA66037-01.

#### 10 A. Field of the Invention

The present invention relates to the field of gene expression, generally, and more specifically it adduces the expression recombinant transgenes in host cells. The invention may be exploited in the production of recombinant proteins *in vitro* or in gene therapy *in vivo*.

#### B. Related Art

The ability to express foreign genes in host cells has become a pivotal tool in molecular biology. For example, expressing proteins in host cells *in vitro* can lead to the large scale production of the protein for use in research or therapy. Examples of proteins that could be used in this manner are hormones, such as insulin, or cytokines, such as the interleukins. Scientists are constantly seeking ways to maximize expression of transgenes when they are imported into host cells.

25

Another important technology affected by foreign gene expression is gene therapy. Attaining high level expression in specific target cells is a key aspect of gene therapy, too. Numerous parameters have been varied in an effort to achieve higher levels of expression including the mode of gene transduction, the vector, the promoter, as well as the dose and routes of administration.

30

Recently Son & Huang (1994) reported that exposure of CDDP-resistant ovarian carcinoma cells to CDDP prior to liposome-mediated gene transfer resulted in enhanced gene transduction. This study utilized an ovarian cancer cell line, 2008, that rapidly acquire CDDP-resistance following exposure. The cells in this study were exposed to CDDP for four to six weeks prior to gene transfer (*in vitro*) or exposed once, one week prior to gene transfer (*in vivo*). According to the authors, their data only indicate that CDDP-resistant cells show improved gene transduction. Thus, from this study, it is unclear whether CDDP-sensitive cells would provide the same results. It also is unclear whether the effect was tied to liposomal transfection methods, or could be more broadly applied.

A similar phenomenon has been observed in primary human foreskin fibroblasts infected with adeno-associated virus following brief exposure to high concentrations of CDDP or other DNA-damaging agents (Alexander *et al.*, 1994; Russell *et al.*, 1995). In this system, these non-malignant cells were rendered more susceptible to transduction by sublethal, but relatively high levels of DNA damaging agents. The treatments were conducted for 16 to 20 hours, followed immediately by transduction. The cause of the increased expression was not correlated with other transduction methodologies. Moreover, it was unclear from the data whether the treatment rendered the cells more competent for AAV transduction, whether increased uptake of AAV was induced or whether increased expression of AAV occurred following internalization.

Thus, it is clear that the prior art does not provide a clear picture, with respect to the effect of DNA-damaging agents, of their effects on the transduction and expression of transgenes in various host cells. There remains a need for a better understanding of these phenomena and for increasing the expression of transgenes in transduced cells.

## SUMMARY OF THE INVENTION

It is, therefore, an object of the present invention to provide improved methods for expression of transgenes in host cells. This applies both to the *in vitro* and *in vivo* applications and includes a variety of different gene delivery systems, DNA-damaging agents, transgenes and target cells. More specifically, the invention provides for

the enhancement of gene expression by providing a DNA damaging agent, preferably to a dividing cell, prior to administration of a vector containing a gene or genes of interest.

In one embodiment, the invention provides a method for enhancing the expression of a transgene comprising (a) contacting a target cell with a DNA-damaging agent; (b) removing the DNA-damaging agent from the target cell; and (c) transferring the transgene into the target cell between about 1-3 days after removing the DNA-damaging agent. More preferably, the transfer of the gene occurs at about 2 days after removal. The target cell preferably is a dividing cell, and more preferably is a tumor cell.

The cell may be drug sensitive or drug insensitive.

The DNA-damaging agent is selected from the group consisting of cisplatin, carboplatin, VP16, teniposide, daunorubicin, doxorubicin, dactinomycin, mitomycin, plicamycin, bleomycin, procarbazine, nitrosourea, cyclophosphamide, bisulfan, melphalan, chlorambucil, ifosfamide, merchlorhehtamine, and ionizing radiation.

Transfer of the transgene is accomplished by a technique selected from the group consisting of liposome-mediated transfection, receptor-mediated internalization and viral infection.

The transgene may be a tumor suppressor, such as p53. The transgene may be under the transcriptional control of a promoter, for example, the CMV IE promoter. Further, the transgene may have, in operable relation thereto, a polyadenylation signal, for example, the SV40 polyadenylation signal. The transgene may be carried in an adenoviral vector.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications

within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

### BRIEF DESCRIPTION OF THE DRAWINGS

5           The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

10   FIG. 1A - Time Course Analysis of CDDP Treatment. Time course of CDDP-induced enhancement of  $\beta$ -gal gene expression in H1299 (open bar) and H460 (hatched bar) cells *in vitro*. Maximal enhancement of  $\beta$ -gal gene expression was observed 2 days after exposure to CDDP (\* =  $p < 0.001$  by ANOVA and Student's t test). Daily infection of untreated H1299 and H460 cells indicated that the transduction efficiency remained  
15   unchanged over the study period; mean  $\pm$  SD (day 0; immediately after removal of CDDP) (n = 6; results of 2 experiments with triplicate samples). Cells were stained for  $\beta$ -gal expression 24 hrs after infection.

FIG. 1B - Dose Response Analysis of CDDP Treatment. Dose response curve of CDDP-  
20   induced enhancement of  $\beta$ -gal gene expression in H1299 (open bar) and H460 (hatched bar) cells. Maximal enhancement of  $\beta$ -gal gene expression was observed in cells exposed to 0.062  $\mu$ g/ml x 24 hrs (\* =  $p < 0.001$  by ANOVA and Student's t test); mean  $\pm$  SD (n = 6; results of two experiments with triplicate samples). Cells were stained for  $\beta$ -gal expression, 24 hrs after infection.

25

FIG. 2A - Enhancement of  $\beta$ -gal Gene Expression in CDDP-Treated Cells in Relation to Increasing MOI's. The  $\beta$ -gal activity of CDDP-treated H1299 cells was quantitated by the ONPG assay, 24 hrs after incubation with Ad/CMV/ $\beta$ -gal. Open circles represent enhancement index; filled circles represent  $\beta$ -gal activity; mean  $\pm$  (n = 4; results of two  
30   experiments with duplicate samples).

FIG. 2B - Enhancement of  $\beta$ -gal Gene Expression in CDDP-Treated Cells in Relation to Increasing MOI's. The  $\beta$ -gal activity of CDDP-treated H460 cells was quantitated by the ONPG assay, 24 hrs after incubation with Ad/CMV/ $\beta$ -gal. Open circles represent enhancement index; filled circles represent  $\beta$ -gal activity; mean  $\pm$  (n = 4; results of two experiments with duplicate samples).

FIG. 3 - Enhancement of  $\beta$ -gal Gene Expression Following Exposure to DNA-Damaging Agents. Enhancement of  $\beta$ -gal expression was observed following treatment with DNA-damaging agents (CDDP, Etoposid (VP16) or ionizing radiation (XRT)), but not to other anti-neoplastic agents (Methotrexate (MTX), Vincristine (VIN), 5-Fluorouracil (5FU) or Transplatinum (TRANS)). Only data of cells treated with drugs (hatched bar) at the concentration of 0.062  $\mu$ g/ml or irradiated with 8 Greys are shown here along with data of non-treated cells (open bar); mean  $\pm$  SD (n = 6; results of two experiments with triplicate samples). Cells were stained for  $\beta$ -gal expression, 24 hrs after infection (\* = p < 0.001 by ANOVA and students t test).

FIG. 4A - Effect of CDDP on Reporter Gene Expression Using Different Vectors. Ad/CMV/ $\beta$ -gal (open bar), Ad/PLL/DNA complex (hatched bar) or lipofectamine (solid bar) carrying a  $\beta$ -gal expressing plasmid were incubated with CDDP-treated (0.062  $\mu$ g/ml for 24 hrs) H1299 cells 48 hrs after CDDP removal; mean  $\pm$  SD (n = 6; results of two experiments, with triplicate samples) (\* = p < 0.001 by ANOVA and Student's t test). Cells were stained for  $\beta$ -gal expression, 24 hrs after vector administration.

FIG. 4B - Effect of CDDP on Reporter Gene Expression Using Different Cell Types. Ad/CMV/ $\beta$ -gal was used to infect malignant and normal cells that were CDDP-treated (hatched bar) or untreated (open bar) with Ad/CMV/ $\beta$ -gal. Infection was performed on day 2 following exposure to CDDP (0.062  $\mu$ g/ml for 24 hrs) or mock exposure with PBS. Only data of cells treated with 0.062  $\mu$ g/ml CDDP are shown. MOI = 1 (H1299; SiHa); MOI = 5 (H460, NHBE); MOI = 10 (H358, H226br, A549). Mean  $\pm$  SD (n = 6; results of two experiments, with triplicate samples). Cells were stained for  $\beta$ -gal expression, 24 hrs after infection.

FIG. 5 - In Vivo Enhancement of  $\beta$ -gal Expression by Sequential Administration of CDDP and Intratumoral Injection of Ad/CMV/ $\beta$ -gal. *In vivo* enhancement of  $\beta$ -gal expression by sequential administration of CDDP (5  $\mu$ g/g body weight; intraperitoneal injection) and intratumoral injection of Ad/CMV/ $\beta$ -gal ( $5 \times 10^8$  viral particles) on the same day (day 0),  
 5 2, 4 or 6 days after CDDP administration; mean  $\pm$  (n = 4 animals) (\* = p < 0.001 by ANOVA and Student's t test). Open bar = CDDP treated animals; hatched bar = PBS treated animals.  $\beta$ -gal analysis was done 24 hrs after vector administration.

FIG. 6 - Enhancement of tumor killing effect by the combination of CDDP and Ad/CMV/p53 *in vitro*. Cells were exposed with CDDP (0.062  $\mu$ g/ml) for 24 hrs, then  
 10 transfected with either Ad/CMV/p53 or dl312 (MOI = 5) 2 days later. Unexposed cells served as controls; \* = p < 0.001 by Student's t test, n = 5.

FIG. 7A - Superior *in vivo* tumor suppression of sequential intraperitoneal CDDP and intratumoral Ad/CMV/p53 administrations. A divided dose regimen ( $1.5 \times 10^{10}$  viral  
 15 particles in three equal doses injected on alternate days) for the administration of CDDP (solid arrow) and Ad/CMV/p53 (open arrow), or a single dose regimen ( $1.5 \times 10^{10}$  viral particles in a single dose) with or without CDDP is shown (p < 0.001 by Student's t test).

FIG. 7B - Superior *in vivo* tumor suppression of sequential intraperitoneal CDDP and intratumoral Ad/CMV/p53 administrations. A single dose regimen with CDDP given  
 20 either before (CDDP on day 0, Ad/CMV/p53 on day 3), concurrent with (CDDP + Ad/CMV/p53 on day 0) or after (Ad/CMV/p53 on day 0, CDDP on day 3) is shown (p < 0.001 by ANOVA and Student's t test; n = 5 per group).

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relies on the observation that treatment of neoplastic cells with DNA-damaging agents, prior to transduction with a transgene, results in the enhanced expression of the transgene. This effect is not observed when the cell is not neoplastic,  
 30 *i.e.*, when the cell exhibits normal growth control. Similarly, the effect is not observed when non-DNA-damaging anti-neoplastic agents are used. However, the effect does not



appear to be limited to particular transgenes nor is it limited to particular transduction methodology, particular neoplastic cells or particular DNA-damaging agents. The invention is described in detail in the following sections.

5 A. Sequential Administration of a DNA-Damaging Agent and a Transgene

In accordance with the present invention, the methods provided for the enhancement of transgene expression are both time and order dependent. Each of these phenomena are discussed below and illustrated in the examples. These data should not be  
10 construed as indicating that time frames or orders outside the preferred embodiments of the present invention are inoperable; to the contrary, data show that simultaneous administration of the p53 tumor suppressor gene in an adenoviral vector and cisplatin results in improved killing of tumor cells, when compared to treatment with the p53  
15 adenoviral vector alone. It is the present inventors' observation, however, that using a particular order and using particular timing, transgene expression may be enhanced over that observed with other protocols.

As the data of the instant examples show, the prior treatment of host cells with a DNA-damaging, followed by provision of a transgene, results in improved expression of  
20 the transgene when compared to simultaneous or subsequent treatment with a DNA-damaging agent, or no DNA-damaging agent at all. In addition, there are particular time frames in which this effect is maximized. Preferably, the *in vitro* administration of the DNA-damaging agent precedes the transduction of the host cell by about 1-3 days (about 24-72 hrs) and, more preferably, about 2 days (about 48 hrs) after removal of the agent. *In*  
25 *vivo*, the time frame may be delayed somewhat, depending on the type and route of administration. Thus, it is suspected that a systemic administration of a chemotherapeutic should precede provision of the transgene by 2-4 days (about 48-96 hrs) and, more preferably, about 3 days (72 hrs).

30 Some minor variation in the optimal treatment times is expected depending on the tumor cell type, the particular transgene, the route of administration, the DNA-damaging agent or the delivery vector. To the extent that the skilled artisan desires, the optimal

timing may be ascertained by performing time course experiments like those set out below in the examples to determine what times will provide the best expression in a given system. It is expected, however, that most will fall within the time frames set forth above.

## 5 B. DNA Damaging Agents

As stated above, the present invention relies on the ability of DNA-damaging agents to facilitate enhanced transgene expression in host cells. For the purposes of this application, DNA-damaging agents are defined as those agents that cause structural  
10 changes in DNA strands existing in the target cell at the time of. The following agents are provided as exemplary of DNA-damaging agents, as defined herein: cisplatin (CDDP), carboplatin, procarbazine, mechlorethamine, cyclophosphamide, ifosfamide, melphalan, chlorambucil, bisulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), and ionizing radiation. Other agents may be  
15 included as long as the adhere to the definition of DNA-damaging agents, provided above.

The following agents are not classified as DNA-damaging agents according to the present invention: transplatinum, 5-fluorouracil, vincristin, vinblastin and methotrexate.

20 Cisplatin (cis-dichlorodimminocisplatin; Mw = 300.05) is a heavy metal compound used as a cancer chemotherapeutic. It has been applied to the treatment of testicular, ovarian, bladder and head/neck cancers. CDDP causes intra- and inter-strand crosslinking of DNA by forming guanine adducts and eventually DNA strand breaks (Eastman, 1990). CDDP is usually administered in 6-8 h periods with 1 liter of fluid and  
25 25 to 50 mg of mannitol. The total dose administered usually is 40-120 mg/m<sup>2</sup> per therapy cycle, but depends on the frequency of cycles and individual tolerance. A common schedule is 20 mg/m<sup>2</sup>/day for five days.

Carboplatin has a mechanism of action and spectrum of clinical activity similar to  
30 CDDP. It is often employed in treating recurrent ovarian cancer, including patients who have previously received cisplatin. Administration is an intravenous infusion over at least 15 min. The usual dose is 360 mg/m<sup>2</sup>, given once every four weeks.

Procarbazine, a methyl hydrazine derivative, is often used clinically against Hodgkin's disease, and also has shown activity against brain and lung tumors, non-Hodgkin's lymphoma, melanoma and myeloma. This drug induces chromosomal damage and chromatid breaks and translocations. The recommended oral daily dose for adults is 100 mg/m<sup>2</sup> for ten days in combination regimens.

Mechlorethamine is a nitrogen mustard that is used in treating Hodgkin's and other lymphomas. Among the effects observed with various nitrogen mustard agents is the opening of imidazole rings or depurination of bases in DNA, both of which can result in serious damage to the DNA strand. The course of therapy consists of the injection of a total dose of 0.4 mg/kg or 10 mg/m<sup>2</sup>. This total dose may be given in either two or four daily consecutive injections, a single administration is preferable. Direct intracavitary administration (0.2 to 0.4 mg/kg) for malignant effusions, particularly of pleural origin, is performed.

Cyclophosphamide, another nitrogen mustard, also alkylates DNA and results in damage to the strands. This drug has been used against Hodgkin's lymphoma, Burkitt's lymphoma and lymphoblastic leukemia. It can be administered orally, intravenously, intramuscularly, intrapleurally and intraperitoneally. A conservative daily dose of 2-3 mg/kg, orally or intravenously, is recommended for patients with more susceptible neoplasms such as lymphomas or leukemias. A higher dosage (4-8 mg/kg daily for six days followed by an oral maintenance dose of 1-5 mg/kg daily) has been used for the treatment of carcinomas and other more resistant neoplasms.

Melphalan is a phenylalanine derivative of nitrogen mustard. It has been used in the treatment of multiple myeloma, malignant melanoma and in carcinoma of the breast and ovary. The usual dose for multiple myeloma is 6 mg daily for a period of 2 to 3 weeks.

Ifosfamide is an analog of cyclophosphamide. It is currently approved for use in combination with other drugs for treatment of germ cell testicular cancer, and clinical

5 trials have shown activity against carcinomas of the lung and cervix, Hodgkin's and non-Hodgkin's lymphomas, and certain sarcomas. The drug usually is infused intravenously over 30 min at a dose of  $1.2 \text{ g/m}^2$  per day for five days. Patients should also receive at least 2 l of oral or intravenous fluid daily. Treatment cycles are usually repeated every three weeks.

10 Chlorambucil is an alkylating agent that has been used against chronic lymphocytic leukemia and primary macroglobulinemia. The standard initial daily dosage is 0.1 to 0.2 mg/kg, continued for at least 3 to 6 weeks. The total daily dose, usually 4 to 10 mg, is given at one time. Bisulfan is another alkylating agent that is used to treat chronic granulocytic leukemia.

15 Nitrosourea is clinically active against lymphomas, malignant melanomas, brain neoplasms and gastrointestinal carcinomas. Chemical decomposition yields reactive intermediates that form single-strand adducts with DNA and then, through a dehalogenation event, forms a second reactive site and cross-links DNA.

20 Dactinomycin is a antibiotic isolated from *Streptomyces*. In addition to blocking transcription, it induces single-strand breaks in DNA, possibly through a free-radical intermediate or as a result of topoisomerase II. It has been used against Hodgkin's and non-Hodgkin's lymphomas. The typical daily dose is 10 to 15  $\mu\text{g/kg/day}$  intravenously for five days. Additional courses may be given at 3- to 4-week intervals in the absence of toxicity. Daily injections of 100 to 400  $\mu\text{g}$  have been given to children for up to fourteen days. Weekly maintenance doses of 7.5  $\mu\text{g/kg}$  have been used.

25 Daunorubicin is an anthracycline antibiotic that, among other actions, induces single- and double-stranded breaks. It is used against acute lymphocytic and acute granulocytic leukemias, acute non-lymphoblastic leukemias and certain lymphomas. The recommended dosage is 30 to 60  $\text{mg/m}^2$  daily for three days.

30 Doxorubicin is a hydroxy analog of daunorubicin, with activity against acute leukemias and malignant lymphomas; it also is active against a number of solid tumors,

including breast cancer. The recommended dose is 60 to 75 mg/m<sup>2</sup>, administered as a single intravenous infusion, repeated after 21 days.

Bleomycins are an important group of antitumor agents derived from *Streptomyces verticillus*. The currently used drug is a mixture of copper-chelating glycopeptides that consist primarily of two closely related agents, bleomycin A<sub>2</sub> and bleomycin B<sub>2</sub>. They show a wide range of activity against a variety of tumors, including squamous carcinomas of the skin, head, neck and lungs, as well as against lymphomas and testicular tumors. Administration is a bolus dose of 15 units/m<sup>2</sup>, twice a day for five days.

Plicamycin is a cytotoxic antibiotic isolated from cultures of *Streptomyces tanashiensis*. It has been used against advanced embryonal tumors of the testes. The recommended dosage for treatment of testicular tumors is 25 to 30 µg/kg daily or on alternate days for eight to ten doses. It usually is administered via intravenous infusion over 4 to 6 hours.

Mitomycin is an antibiotic isolated from *Streptomyces caespitosus*. The drug both inhibits DNA synthesis and cross-links DNA; single-strand breaks are induced by the removal of mitomycin.. It is used in the palliative treatment of gastric adenocarcinoma, in conjunction with 5-FU and doxorubicin. It also has produced temporary benefits in carcinomas of the cervix, colon, rectum, pancreas, breast, bladder, head and neck, and lung, and in melanomas.

Etoposide (Mw = 588.58) is a semi-synthetic derivative of podophyllotoxin with antineoplastic activity against testicular tumors and small-cell carcinoma of the lung, usually in combination with cisplatin. It also is active against non-Hodgkin's lymphomas, acute non-lymphocytic leukemia, carcinoma of the breast and Kaposi's sarcoma. Etoposide forms a tertiary complex between DNA and topoisomerase II and eventually produces protein-linked DNA single and double strand breaks (Kaufman, 1989). Intravenous dosage for testicular tumors is 50-100 mg/m<sup>2</sup> for five days, or 100 mg/m<sup>2</sup> on alternate days for three doses. For small-cell carcinoma, the dose is 35 mg/m<sup>2</sup> daily for

four days to 50 mg/m<sup>2</sup> daily for five days. If given orally, the dosage should be doubled. Cycles of therapy are usually repeated every three to four weeks.

5 Teniposide is a thiophene derivative of etoposide that is used in the treatment of refractory acute lymphoblastic leukemia in children. Administration is via intravenous infusion in doses that range from 50 mg/m<sup>2</sup> per day for five days to 165 mg/m<sup>2</sup> per day, twice weekly.

10 Radiation is an important tool in tumor therapy. Radiation falls into two major categories - electromagnetic radiation (waves of varying frequency, *e.g.*, x-rays) and subatomic particle radiation (alpha, beta (electron), neutron, proton, meson and heavy ion). Gamma emissions are a form of electromagnetic radiation emitted from radioactive isotopes of radium, cobalt and other elements.

15 Radiation therapy transfers discrete energy units, called photons, to tissues causing damage to both normal and malignant cells. Ionizing irradiation stimulates production of oxygen free radicals which react with macromolecules and induces DNA damage (Cole *et al.*, 1980). "Early" radiation effects include damage to proliferating cells, while "late" effects involve cell death and affect many different kinds of cells. Fortunately, radiation  
20 exploits the differential effects on malignant *versus* non-malignant cells, namely, that rapidly proliferating cells undergoing significant DNA synthesis suffer more severe effects from the DNA damage induced by radiation.

The dose of radiation is dependent upon tissue and tumor type. The treatment is  
25 usually fractionated to prevent toxicity and can be in the 1-5 Grey range over a several week period. For the treatment of rectal cancer radiation, 45 Grey total dose is given (1.8 Grey dose/day, Monday through Friday). For the enhancement of gene expression, a single dose of between 1 and 8 Grey is contemplated for gamma radiation.

### C. Transgenes and Expression Constructs

The transgenes of the present invention may encode any protein of interest.

Various proteins are useful for their biological activities *in vitro* and *in vivo*. For example, cytokines and hormones have already found use in treating certain diseases. For example, the interleukins (IL-1 to 11), interferon, human growth hormone, insulin, insulin-like growth factor, prolactin, placental lactogen, luteinizing hormone, follicle stimulating hormone, chorionic gonadotropic, thyroid-stimulating hormone, glucagon, somatostatin, calcitonin, vasopressin, vasostatin, vasotocin, gastrin, amylin, growth hormone releasing factor, growth hormone releasing hormone, luteinizing hormone releasing hormone, thymidine kinase, interleukin-1-beta converting enzyme and others are suitable subjects of the present invention.

In addition, the use of tumor suppressor genes such as p53, C-CAM, retinoblastoma gene, herpesvirus and p16 are candidates for gene transfer. These proteins act to regulate cell growth, and their reduction, mutation or deletion can affect the ability of cells to undergo normal cell senescence or apoptosis. Introducing transgenes encoding these proteins can reestablish normal cell growth, and expression of high levels of these proteins can even overcome defects in other genes that cause neoplastic proliferation.

20

p53 is a 53 kD nuclear phosphoprotein of 375 amino acids that controls cell proliferation. Mutations to the p53 gene, and allele loss on chromosome 17p, where this gene is located, are among the most frequent alterations identified in human malignancies. The p53 protein is highly conserved through evolution and is expressed in most normal tissues. Wild-type p53 has been shown to be involved in control of the cell cycle (Mercer, 1992), transcriptional regulation (Fields *et al.*, 1990; Mietz *et al.*, 1992), DNA replication (Wilcock and Lane, 1991; Bargonetti *et al.*, 1991) and induction of apoptosis (Yonish-Rouach *et al.*, 1991; Shaw *et al.*, 1992).

25

Various mutant p53 alleles are known in which a single base substitution results in the synthesis of proteins that have quite different growth regulatory properties and, ultimately, lead to malignancies (Hollstein *et al.*, 1991). In fact, the p53 gene has been

30

found to be the most frequently mutated gene in common human cancers (Hollstein *et al.*, 1991; Weinberg, 1991), and is particularly associated with those cancers linked to cigarette smoke (Hollstein *et al.*, 1991; Zakut-Houri *et al.*, 1985). The overexpression of p53 in breast tumors has also been documented (Casey *et al.*, 1991).

5

Unlike other oncogenes, however, p53 point mutations are known to occur in at least 30 distinct codons, often creating dominant alleles that produce shifts in cell phenotype without a reduction to homozygosity. Additionally, many of these dominant negative alleles appear to be tolerated in the organism and passed on in the germ line. Various mutant alleles appear to range from minimally dysfunctional to strongly penetrant, dominant negative alleles (Weinberg, 1991).

10

Casey and colleagues have reported that transfection of DNA encoding wild-type p53 into two human breast cancer cell lines restores growth suppression control in such cells (Casey *et al.*, 1991). A similar effect has also been demonstrated on transfection of wild-type, but not mutant, p53 into human lung cancer cell lines (Takahasi *et al.*, 1992). The p53 appears dominant over the mutant gene and will select against proliferation when transfected into cells with the mutant gene. Normal expression of the transfected p53 does not affect the growth of cells with endogenous p53.

20

Alternatively, the transgenes may encode antisense oligonucleotides that hybridize, under intracellular conditions, to a target nucleic acid. The target nucleic acid may be a DNA molecule or an RNA molecule. Hybridization results in the inhibition of transcription and or translation of the protein encoded by the target nucleic acid. The design of antisense constructs, based on the sequence of genes, will be evident to those of skill in the art.

25

The term "antisense nucleic acid" is intended to refer to the oligonucleotides complementary to the base sequences of a target DNA or RNA. Antisense oligonucleotides, when introduced into a target cell, specifically bind to their target nucleic acid and interfere with transcription, RNA processing, transport, translation and/or stability. Targeting double-stranded (ds) DNA with oligos or oligonucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation.

30



Antisense constructs may be designed to bind to the promoter and other control regions, exons, introns or even exon-intron boundaries of a gene. Antisense RNA constructs, or DNA encoding such antisense RNA's, may be employed to inhibit gene transcription or translation or both within a host cell, either *in vitro* or *in vivo*, such as within a host animal, including a human subject. Nucleic acid sequences which comprise "complementary nucleotides" are those which are capable of base-pairing according to the standard Watson-Crick complementarity rules. That is, that the larger purines will base pair with the smaller pyrimidines to form combinations of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T), in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. Inclusion of less common bases such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others in hybridizing sequences does not interfere with pairing.

As used herein, the terms "complementary" or "antisense sequences" mean nucleic acid sequences that are substantially complementary over their entire length and have very few base mismatches. For example, nucleic acid sequences of fifteen bases in length may be termed complementary when they have a complementary nucleotide at thirteen or fourteen positions with only a single mismatch. Naturally, nucleic acid sequences which are "completely complementary" will be nucleic acid sequences which are entirely complementary throughout their entire length and have no base mismatches.

Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology, but also contains a non-homologous region (*e.g.*, a ribozyme) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

While all or part of the gene sequence may be employed in the context of antisense construction, statistically, any sequence of 17 bases long should occur only once in the human genome and, therefore, suffice to specify a unique target sequence. Although shorter oligomers are easier to make and increase *in vivo* accessibility, numerous other factors are

involved in determining the specificity of hybridization. Both binding affinity and sequence specificity of an oligonucleotide to its complementary target increases with increasing length. It is contemplated that oligonucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more base pairs will be used. One can readily determine whether a given antisense nucleic acid is effective at targeting of the corresponding host cell gene simply by testing the constructs *in vitro* to determine whether the endogenous gene's function is affected or whether the expression of related genes having complementary sequences is affected.

In certain embodiments, one may wish to employ antisense constructs which include other elements, for example, those which include C-5 propyne pyrimidines. Oligonucleotides which contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression (Wagner *et al.*, 1993).

As an alternative to targeted antisense delivery, targeted ribozymes may be used. The term "ribozyme" is refers to an RNA-based enzyme capable of targeting and cleaving particular base sequences in target DNA and RNA. Ribozymes can either be targeted directly to cells, in the form of RNA oligonucleotides incorporating ribozyme sequences, or introduced into the cell as an expression construct encoding the desired ribozymal RNA. Ribozymes may be used and applied in much the same way as described for antisense nucleic acids. Ribozyme sequences also may be modified in much the same way as described for antisense nucleic acids. For example, one could incorporate non-Watson-Crick bases, or make mixed RNA/DNA oligonucleotides, or modify the phosphodiester backbone.

The term "cDNA" is intended to refer to DNA prepared using messenger RNA (mRNA) as template. The advantage of using a cDNA, as opposed to genomic DNA or DNA polymerized from a genomic, non- or partially-processed RNA template, is that the cDNA does not contain any non-coding sequences but, rather, contains only the coding region of the corresponding protein. There may be times when the full or partial genomic sequence is preferred, such as where the non-coding regions are required for optimal expression or where non-coding regions such as introns are to be targeted in an antisense strategy.

Throughout this application, the term "expression construct" is meant to include any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a protein, but it need not be. Thus, in certain embodiments, expression includes both transcription of a gene and translation of a RNA into a gene product. In other embodiments, expression only includes transcription of the nucleic acid, *i.e.*, antisense and ribozymes.

In preferred embodiments, the nucleic acid is under transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrase "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

The term promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (tk) and SV40 early transcription units. These studies, augmented by more recent work, have shown that promoters are composed of discrete functional modules, each consisting of approximately 7-20 bp of DNA, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

The particular promoter that is employed to control the expression of a nucleic acid is not believed to be critical, so long as it is capable of expressing the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a human or viral promoter.

In various embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter and the Rous sarcoma virus long terminal repeat can be used to obtain high-level expression of transgenes. The use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression of a transgene is contemplated as well, provided that the levels of expression are sufficient for a given purpose.

By employing a promoter with well-known properties, the level and pattern of expression following transfection can be optimized. For example, selection of a promoter which is active specifically in lung cells, such as tyrosinase (melanoma), alpha-fetoprotein and albumin (liver tumors), CC10 (lung tumor) and prostate-specific antigen (prostate tumor) will permit tissue-specific expression. Further, selection of a promoter that is regulated in response to specific physiologic signals can permit inducible expression. For example, with human PAI-1 promoter, expression is inducible by tumor necrosis factor. Tables 1 and 2 list several elements/promoters which may be employed, in the context of the present invention,

to regulate the expression of a transgene. This list is not intended to be exhaustive of all the possible elements involved in the promotion of transgene expression but, merely, to be exemplary thereof.

5 Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of DNA with enhancer activity are organized much like promoters. That is, they are composed of many individual elements,  
10 each of which binds to one or more transcriptional proteins.

The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have  
15 one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

20 Below is a list of viral promoters, cellular promoters/enhancers and inducible promoters/enhancers that could be used in combination with the nucleic acid encoding a transgene in an expression construct (Table 1 and Table 2). Additionally any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base EPDB) could also be used to drive expression of a transgene. Use of a T3, T7 or SP6 cytoplasmic  
25 expression system is another possible embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

TABLE 1

PROMOTER
Immunoglobulin Heavy Chain
Immunoglobulin Light Chain
T-Cell Receptor
HLA DQ $\alpha$ and DQ $\beta$
$\beta$ -Interferon
Interleukin-2
Interleukin-2 Receptor
MHC Class II 5
MHC Class II HLA-DR $\alpha$
$\beta$ -Actin
Muscle Creatine Kinase
Prealbumin (Transthyretin)
Elastase I
Metallothionein
Collagenase
Albumin Gene
$\alpha$ -Fetoprotein
$\tau$ -Globin
$\beta$ -Globin
c-fos

TABLE 1 CONT'D

PROMOTER
c-HA-ras
Insulin
Neural Cell Adhesion Molecule (NCAM)
$\alpha$ 1-Antitrypsin
H2B (TH2B) Histone
Mouse or Type I Collagen
Glucose-Regulated Proteins (GRP94 and GRP78)
Rat Growth Hormone
Human Serum Amyloid A (SAA)
Troponin I (TN I)
Platelet-Derived Growth Factor
Duchenne Muscular Dystrophy
SV40
Polyoma
Retroviruses
Papilloma Virus
Hepatitis B Virus
Human Immunodeficiency Virus
Cytomegalovirus
Gibbon Ape Leukemia Virus

TABLE 2

Element	Inducer
MT II	Phorbol Ester (TFA) Heavy metals
MMTV (mouse mammary tumor virus)	Glucocorticoids
$\beta$ -Interferon	poly(rI)X poly(rc)
Adenovirus 5 E2	Ela
c-jun	Phorbol Ester (TPA), H <sub>2</sub> O <sub>2</sub>
Collagenase	Phorbol Ester (TPA)
Stromelysin	Phorbol Ester (TPA), IL-1
SV40	Phorbol Ester (TFA)
Murine MX Gene	Interferon, Newcastle Disease Virus
GRP78 Gene	A23187
$\alpha$ -2-Macroglobulin	IL-6
Vimentin	Serum
MHC Class I Gene H-2kB	Interferon
HSP70	Ela, SV40 Large T Antigen
Proliferin	Phorbol Ester-TPA
Tumor Necrosis Factor	FMA
Thyroid Stimulating Hormone $\alpha$ Gene	Thyroid Hormone



Where a cDNA insert is employed, one will typically desire to include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal and the bovine growth hormone polyadenylation signal, convenient and known to function well in various target cells. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

Specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements (Bittner *et al.*, 1987).

In preferred embodiments of the invention, the expression construct comprises a virus or engineered construct derived from a viral genome. The ability of certain viruses to enter cells via receptor-mediated endocytosis and to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign genes into mammalian cells (Ridgeway, 1988; Nicolas and Rubenstein, 1988; Baichwal and Sugden, 1986; Temin, 1986). The first viruses used as gene vectors were DNA viruses including the papovaviruses (simian virus 40, bovine papilloma virus, and polyoma) (Ridgeway, 1988; Baichwal and Sugden, 1986), adenoviruses (Ridgeway, 1988; Baichwal and Sugden, 1986). These have a relatively low capacity for foreign DNA sequences and have a restricted host spectrum. Furthermore, their oncogenic potential and cytopathic effects in permissive cells raise safety concerns. They can accommodate only up

to 8 kilobases of foreign genetic material but can be readily introduced in a variety of cell lines and laboratory animals (Nicolas and Rubenstein, 1988; Temin, 1986).

(i) Retroviruses

5

The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the gag gene, termed  $\Psi$ , functions as a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome (Coffin, 1990).

15

In order to construct a retroviral vector, a nucleic acid encoding a transgene is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and  $\Psi$  components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a human cDNA, together with the retroviral LTR and  $\Psi$  sequences is introduced into this cell line (by calcium phosphate precipitation for example), the  $\Psi$  sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind *et al.*, 1975).

25

30

A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical

addition of lactose residues to the viral envelope. This modification could permit the specific infection of hepatocytes via sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was designed in which  
5 biotinylated antibodies against a retroviral envelope protein and against a specific cell  
receptor were used. The antibodies were coupled via the biotin components by using  
streptavidin (Roux *et al.*, 1989). Using antibodies against major histocompatibility complex  
class I and class II antigens, they demonstrated the infection of a variety of human cells that  
bore those surface antigens with an ecotropic virus *in vitro* (Roux *et al.*, 1989).

10

There are certain limitations to the use of retrovirus vectors in all aspects of the  
present invention. For example, retrovirus vectors usually integrate into random sites in the  
cell genome. This can lead to insertional mutagenesis through the interruption of host genes  
or through the insertion of viral regulatory sequences that can interfere with the function of  
15 flanking genes (Varmus *et al.*, 1981). Another concern with the use of defective retrovirus  
vectors is the potential appearance of wild-type replication-competent virus in the packaging  
cells. This can result from recombination events in which the intact  $\Psi$  sequence from the  
recombinant virus inserts upstream from the gag, pol, env sequence integrated in the host cell  
genome. However, new packaging cell lines are now available that should greatly decrease  
20 the likelihood of recombination (Markowitz *et al.*, 1988; Hersdorffer *et al.*, 1990).

25

One limitation to the use of retrovirus vectors *in vivo* is the limited ability to produce  
retroviral vector titers greater than  $10^6$  infectious U/mL. Titers 10- to 1,000-fold higher are  
necessary for many *in vivo* applications.

30

## (ii) Adenovirus

Knowledge of the genetic organization of adenovirus, a 36 kB, linear and double-  
stranded DNA virus, allows substitution of a large piece of adenoviral DNA with foreign  
30 sequences up to 7 kB (Grunhaus and Horwitz, 1992). In contrast to retrovirus, the infection  
of adenoviral DNA into host cells does not result in chromosomal integration because  
adenoviral DNA can replicate in an episomal manner without potential genotoxicity. Also,

adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in the human.

5

Adenovirus is particularly suitable for use as a gene transfer vector because of its mid-sized genome, ease of manipulation, high titer, wide target-cell range, and high infectivity. Both ends of the viral genome contain 100-200 base pair (bp) inverted terminal repeats (ITR), which are *cis* elements necessary for viral DNA replication and packaging.

10 The early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late

15 gene expression, and host cell shut off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNAs issued from this promoter possess a 5' tripartite leader (TL) sequence which

20 makes them preferred mRNAs for translation.

In the current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated from

25 this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure. Use of the YAC system is an alternative approach for the production of recombinant adenovirus.

Generation and propagation of the current adenovirus vectors, which are replication

30 deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham, *et al.*, 1977). Since the E3 region is dispensable from the adenovirus

genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the E3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury, *et al.*, 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1 deleted virus is incomplete. For example, leakage of viral gene expression has been observed with the currently available adenovirus vectors at high multiplicities of infection (Mulligan, 1993).

Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, *e.g.*, Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the preferred helper cell line is 293.

Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain the conditional replication-defective adenovirus vector for use in the method of the present invention. This is because Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical adenoviral vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most convenient to introduce the transgene nucleic acid into the position from which the E1 coding sequences have been removed. However, the position of insertion of the coding

region within the adenovirus sequences is not critical to the present invention. The nucleic acid encoding a transcription unit also may be inserted in lieu of the deleted E3 region in E3 replacement vectors as described previously by Karlsson *et al.* (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

5

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titers, *e.g.*,  $10^9$ - $10^{11}$  plaque-forming unit (PFU)/ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal, and therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch *et al.*, 1963; Top *et al.*, 1971), demonstrating their safety and therapeutic potential as *in vivo* gene transfer vectors.

10

Adenovirus vectors have been used in eukaryotic gene expression (Levrero *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet and Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Experiments in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injection (Herz and Gerard, 1993), and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993). In addition to these protocols, the present invention also contemplates direct tumoral injection.

20

### (iii) Other Viral Vectors as Expression Constructs

25

Other viral vectors may be employed as expression constructs in the present invention. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar *et al.*, 1988) adeno-associated virus (AAV) (Ridgeway, 1988; Baichwal and Sugden, 1986; Hermonat and Muzycska, 1984) and herpesviruses may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar *et al.*, 1988; Horwich *et al.*, 1990).

30

With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang *et al.* recently introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B' virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

#### D. Methods of Gene Transfer

In order to effect expression of sense or antisense transgene constructs, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain disease states. As described above, the one mechanism for delivery is via viral infection where the expression construct is encapsidated in an infectious viral particle.

Several non-viral methods for the transfer of expression constructs into cultured mammalian cells also are contemplated by the present invention. These include calcium phosphate precipitation (Graham and Van Der Eb, 1973; Chen and Okayama, 1987; Rippe *et al.*, 1990) DEAE-dextran (Gopal, 1985), electroporation (Tur-Kaspa *et al.*, 1986; Potter *et al.*, 1984), direct microinjection (Harland and Weintraub, 1985), DNA-loaded liposomes (Nicolau and Sene, 1982; Fraley *et al.*, 1979) and lipofectamine-DNA complexes, cell sonication (Fechheimer *et al.*, 1987), gene bombardment using high velocity microprojectiles (Yang *et al.*, 1990), and receptor-mediated transfection (Wu and Wu, 1987; Wu and Wu, 1988). Some of these techniques may be successfully adapted for *in vivo* or *ex vivo* use.

In one embodiment of the invention, the expression construct may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of  $\text{CaPO}_4$  precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Neshif (1986) also demonstrated that direct intraperitoneal injection of  $\text{CaPO}_4$  precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a transgene may also be transferred in a similar manner *in vivo* and express the corresponding protein.

Another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ, *i.e.*, *ex vivo* treatment. Again, DNA encoding a transgene may be delivered via this method and still be incorporated by the present invention.

In a further embodiment of the invention, the expression construct may be entrapped in a liposome. Liposomes are vesicular structures characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-



rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). Also contemplated are lipofectamine-DNA complexes.

5 Liposome-mediated nucleic acid delivery and expression of foreign DNA *in vitro* has been very successful. Wong *et al.* (1980) demonstrated the feasibility of liposome-mediated delivery and expression of foreign DNA in cultured chick embryo, HeLa and hepatoma cells. Nicolau *et al.* (1987) accomplished successful liposome-mediated gene transfer in rats after intravenous injection.

10

In certain embodiments of the invention, the liposome may be complexed with a hemagglutinating virus (HVJ). This has been shown to facilitate fusion with the cell membrane and promote cell entry of liposome-encapsulated DNA (Kaneda *et al.*, 1989). In other embodiments, the liposome may be complexed or employed in conjunction with  
15 nuclear non-histone chromosomal proteins (HMG-1) (Kato *et al.*, 1991). In yet further embodiments, the liposome may be complexed or employed in conjunction with both HVJ and HMG-1. In that such expression constructs have been successfully employed in transfer and expression of nucleic acid *in vitro* and *in vivo*, then they are applicable for the present invention. Where a bacterial promoter is employed in the DNA construct, it also will be  
20 desirable to include within the liposome an appropriate bacterial polymerase.

Other expression constructs which can be employed to deliver a nucleic acid encoding a transgene into cells are receptor-mediated delivery vehicles. These take advantage of the selective uptake of macromolecules by receptor-mediated endocytosis in  
25 almost all eukaryotic cells. Because of the cell type-specific distribution of various receptors, the delivery can be highly specific (Wu and Wu, 1993).

Receptor-mediated gene targeting vehicles generally consist of two components: a cell receptor-specific ligand and a DNA-binding agent. Several ligands have been used for  
30 receptor-mediated gene transfer. The most extensively characterized ligands are asialoorosomucoid (ASOR) (Wu and Wu, 1987) and transferrin (Wagner *et al.*, 1990).

Recently, a synthetic neoglycoprotein, which recognizes the same receptor as ASOR, has been used as a gene delivery vehicle (Ferkol *et al.*, 1993; Perales *et al.*, 1994).

In other embodiments, the delivery vehicle may comprise a ligand and a liposome. For example, Nicolau *et al.* (1987) employed lactosyl-ceramide, a galactose-terminal asialganglioside, incorporated into liposomes and observed an increase in the uptake of the insulin gene by hepatocytes. Thus, it is feasible that a nucleic acid may be specifically delivered into a cell type such as lung, epithelial or tumor cells, by any number of receptor-ligand systems with or without liposomes. For example, epidermal growth factor (EGF) may be used as the receptor for mediated delivery of a nucleic acid encoding a transgene in many tumor cells that exhibit upregulation of EGF receptor. Mannose can be used to target the mannose receptor on liver cells. Also, antibodies to CD5 (CLL), CD22 (lymphoma), CD25 (T-cell leukemia) and MAA (melanoma) can similarly be used as targeting moieties.

In certain embodiments, gene transfer may more easily be performed under *ex vivo* conditions. *Ex vivo* gene therapy refers to the isolation of cells from an animal, the delivery of a nucleic acid into the cells, *in vitro*, and then the return of the modified cells back into an animal. This may involve the surgical removal of tissue/organs from an animal or the primary culture of cells and tissues. Anderson *et al.*, U.S. Patent 5,399,346, and incorporated herein in its entirety, disclose *ex vivo* therapeutic methods.

Primary mammalian cell cultures may be prepared in various ways. In order for the cells to be kept viable while *in vitro* and in contact with the expression construct, it is necessary to ensure that the cells maintain contact with the correct ratio of oxygen and carbon dioxide and nutrients but are protected from microbial contamination. Cell culture techniques are well documented and are disclosed herein by reference (Freshner, 1992).

During *in vitro* culture conditions, the expression construct delivers a nucleic acid to the cells and the protein is expressed. Finally, the cells may be reintroduced into the original animal, or administered into a distinct animal, in a pharmaceutically acceptable form by any of the means described below. Thus, providing an *ex vivo* method of treating a mammal with a pathologic condition is within the scope of the invention.

E. Recombinant Protein Production *In Vitro*

Another embodiment of the present invention involves the use of gene transfer to generate recombinant cells lines *in vitro* for the production of recombinant proteins. The gene of interest may be transferred as described above into appropriate host cells followed by culture of cells under the appropriate conditions. The gene for virtually any polypeptide may be employed in this manner. The generation of recombinant expression vectors, and the elements included therein, are discussed above.

Examples of useful mammalian host cell lines are Vero and HeLa cells and cell lines of Chinese hamster ovary, W138, BHK, COS-7, 293, HepG2, 3T3, RIN and MDCK cells. In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and process the gene product in the manner desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to insure the correct modification and processing of the foreign protein expressed.

A number of selection systems may be used including, but not limited to, the herpes simplex virus thymidine kinase, hypoxanthine-guanine phosphoribosyltransferase and adenine phosphoribosyltransferase genes, in *tk-*, *hgprt-* or *aprt-* cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection for *dhfr*, that confers resistance to; *gpt*, that confers resistance to mycophenolic acid; *neo*, that confers resistance to the aminoglycoside G418; and *hygro*, that confers resistance to hygromycin.

Animal cells can be propagated *in vitro* in two modes: as non-anchorage dependent cells growing in suspension throughout the bulk of the culture or as anchorage-dependent cells requiring attachment to a solid substrate for their propagation (*i.e.*, a monolayer type of cell growth).

Non-anchorage dependent or suspension cultures from continuous established cell lines are the most widely used means of large scale production of cells and cell products. However, suspension cultured cells have limitations, such as tumorigenic potential and lower protein production than adherent cells.

5

Large scale suspension culture of mammalian cells in stirred tanks is a common method for production of recombinant proteins. Two suspension culture reactor designs are in wide use - the stirred reactor and the airlift reactor. The stirred design has successfully been used on an 8000 liter capacity for the production of interferon (Phillips *et al.*, 1985; Mizrahi, 1983). Cells are grown in a stainless steel tank with a height-to-diameter ratio of 1:1 to 3:1. The culture is usually mixed with one or more agitators, based on bladed disks or marine propeller patterns. Agitator systems offering less shear forces than blades have been described. Agitation may be driven either directly or indirectly by magnetically coupled drives. Indirect drives reduce the risk of microbial contamination through seals on stirrer shafts.

15

The airlift reactor, also initially described for microbial fermentation and later adapted for mammalian culture, relies on a gas stream to both mix and oxygenate the culture. The gas stream enters a riser section of the reactor and drives circulation. Gas disengages at the culture surface, causing denser liquid free of gas bubbles to travel downward in the downcomer section of the reactor. The main advantage of this design is the simplicity and lack of need for mechanical mixing. Typically, the height-to-diameter ratio is 10:1. The airlift reactor scales up relatively easily, has good mass transfer of gases and generates relatively low shear forces.

25

#### F. In Vivo Gene Therapy Applications

In another embodiment of the present invention, methods for improved gene therapy are provided. The present invention contemplates the use of gene therapeutic vectors, in conjunction with DNA-damaging agent treatment, to provide for high level expression of transgenes *in vivo*. This will be accomplished by treating an individual with an effective amount of a DNA-damaging agent, followed by administration of a

30

therapeutic gene. As stated above, the timing of the administration is important for achieving maximal enhancement of expression. The 1-3 day time period between administration of the DNA-damaging agent and administration of the transgene for *in vitro* may be delayed somewhat for *in vivo* applications, especially where the DNA-damaging agent is administered systematically. However, where the DNA-damaging agent is given locally to the site of administration of the transgene, the 1-3 window should be appropriate.

Administration of the DNA-damaging agent to patient will follow well-established protocols. For example, the administration of cisplatin would occur via intravenous infusion over about 8 hours with adequate prior hydration to minimize nephrotoxicity. The particular dose is dependent upon tumor type and size and the overall condition of the patient. The dose ranges from 60 mg/m<sup>2</sup> to 100 mg/m<sup>2</sup>. Two to three days following cisplatin treatment, the vector system would be administered, for example, via intratumoral injection. It is anticipated that the treatment cycles would be repeated every four weeks as necessary.

Where clinical application of an expression construct comprising a nucleic acid encoding a transgene is contemplated, it will be necessary to prepare the complex as a pharmaceutical composition appropriate for the intended application. Generally this will entail preparing a pharmaceutical composition that is essentially free of pyrogens, as well as any other impurities that could be harmful to humans or animals. One also will generally desire to employ appropriate salts and buffers to render the complex stable and allow for complex uptake by target cells.

Aqueous compositions of the present invention comprise an effective amount of the expression construct and nucleic acid, dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. Such compositions can also be referred to as inocula. The phrases "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such

media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

5

Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain

10 a preservative to prevent the growth of microorganisms.

15

The expression constructs and delivery vehicles of the present invention may include classic pharmaceutical preparations. Administration of therapeutic compositions according to the present invention will be via any common route so long as the target tissue is available via that route. This includes oral, nasal, buccal, rectal, vaginal or topical. Topical administration would be particularly advantageous for treatment of skin cancers, to prevent chemotherapy-induced alopecia or other dermal hyperproliferative disorder. Alternatively, administration will be by orthotopic, intradermal subcutaneous, intramuscular, intraperitoneal or intravenous injection. Such compositions would normally be administered

20 as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients.

25

The therapeutic compositions of the present invention are advantageously administered in the form of injectable compositions either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. These preparations also may be emulsified. A typical composition for such purpose comprises a pharmaceutically acceptable carrier. For instance, the composition may contain 10 mg, 25 mg, 50 mg or up to about 100 mg of human serum albumin per milliliter of phosphate buffered saline. Other pharmaceutically acceptable carriers include aqueous

30 solutions, non-toxic excipients, including salts, preservatives, buffers and the like. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil and injectable organic esters such as ethyloleate. Aqueous carriers include water.

alcoholic/aqueous solutions, saline solutions, parenteral vehicles such as sodium chloride, Ringer's dextrose, *etc.* Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial agents, anti-oxidants, chelating agents and inert gases. The pH and exact concentration of the various components the pharmaceutical composition  
5 are adjusted according to well known parameters.

Additional formulations are suitable for oral administration. Oral formulations include such typical excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like.  
10 The compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders. When the route is topical, the form may be a cream, ointment, salve or spray.

An effective amount of the therapeutic agent is determined based on the intended  
15 goal, for example, in the case of tumor therapy: (i) inhibition of tumor cell proliferation or (ii) elimination of tumor cells. The term "unit dose" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined-quantity of the therapeutic composition calculated to produce the desired responses, discussed above, in association with its administration, *i.e.*, the appropriate route and treatment regimen. The quantity to be  
20 administered, both according to number of treatments and unit dose, depends on the subject to be treated, the state of the subject and the protection desired. Precise amounts of the therapeutic composition also depend on the judgment of the practitioner and are peculiar to each individual.

## 25 G. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the  
30 inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific

embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

## EXAMPLE 1: ENHANCEMENT OF GENE EXPRESSION IN HUMAN 5 CARCINOMA CELLS BY DNA-DAMAGING AGENTS FOLLOWING GENE TRANSFER

### MATERIALS AND METHODS

#### 10 Cells and Culture Conditions

The human non-small cell lung carcinoma cell lines H1299 and H460 were grown in RPMI-1640 medium supplemented with 5% heat-inactivated fetal calf serum (FCS), 10 mM glutamine, 100 units/ml of penicillin, 100 µg/ml of streptomycin and 0.25 µg/ml of  
15 amphotericin B. Human non-small cell lung carcinoma cell lines H358, H226br, and the cervical cancer cell line SiHa were maintained in complete RPMI-1640 medium supplemented with 10% FCS. The cell line A549 (human, non-small cell lung carcinoma) was grown in Ham's F12 nutrient mixture solution (Gibco BRL, Gaithersburg, MD) similarly supplemented with antibiotics, glutamine and 10% FCS. Primary normal human  
20 bronchial epithelial (NHBE) cells (Clonetics Corporation, San Diego, CA), were grown in serum-free optimized growth medium and subcultured under conditions suggested by the manufacturer.

#### *In Vitro* Characterization of the Effects of CDDP Exposure on β-gal Expression

25

**Gene Delivery Protocol:** *In vitro* gene delivery was performed by incubating cells in six-well plates (Falcon Plastics, Lincoln Park, NJ) for two hours with Adv/CMV/β-gal in appropriate medium supplemented with 2% FCS. The multiplicity of infection (MOI: number of viral particles per target cell) was based on cell counts of untreated wells.  
30 Fresh, complete medium with the appropriate concentration of FCS was then added to the wells at the end of the infection period and cells were then incubated for 20 hours at 37°C,



5% CO<sub>2</sub>. After washing the cells with ice cold PBS, the cells were fixed with ice-cold 1.25% glutaraldehyde and stained with X-gal (5-bromo-4-chloro-3-indolyl-β-d-galactoside; Gibco BRL) as previously described (MacGregor *et al.*, 1987). The transduction efficiency by Adv/CMV/β-gal was determined by the percentage of positive stained cells (blue) for β-galactosidase activity (1000 cells counted per well).

**Time Course and Dose Response:** To determine the time course of enhancement, H1299 and H460 cells were seeded in 6-well plates ( $2 \times 10^5$  cells/2 mls/well). After an overnight incubation period (37° C, 5% CO<sub>2</sub>), the initial medium was replaced with freshly prepared media (2 ml) containing 0.062 μg/ml CDDP (Sigma Chemicals, St. Louis, MO) (CDDP was prepared fresh by dissolving in distilled water) and the cells were then incubated for 24 hours at 37°C, with 5% CO<sub>2</sub>. CDDP-treated cells were then washed twice with PBS and subsequently infected with Ad/CMV/β-gal (MOI = 1 for H1299 and MOI = 5 for H460) on days 0, 1, 2, 3, 4, 5 or 6 following CDDP removal, to determine the time point at which maximal enhancement of expression was occurring. To establish the dose of CDDP needed to produce maximum enhancement, the cells were treated with different concentrations of CDDP (serial, 1:4 dilutions, ranging from 4 μg/ml to 0.0002 μg/ml, 5 ml per 60 mm dish) for 24 hours and subsequently incubated with Ad/CMV/β-gal 48 hours after removal of CDDP. To study if exposure of malignant cells to different classes of antineoplastic agents would lead to a similar elevation of gene expression, H1299 cells were treated with vincristine (Eli Lilly and Co, Indianapolis, IN), methotrexate (Lederle Pharmaceuticals, Pearl River, NY), 5-fluorouracil (SoloPak Laboratories, Elk Grove Village, IL), epotoside (Bristol Laboratories, Princeton, NJ) and transplatin (the isomer of CDDP that has no anticancer activity, Sigma Chemical, St. Louis, MO) in the similar fashion as CDDP treatment and then infected with Ad/CMV/β-gal 48 hours after removal of drug-containing media. H1299 cells were treated with 2, 4, 8, or 16 Grey (Gy) of ionizing irradiation and then similarly infected 72 hours after irradiation. Untreated cells that were similarly infected, served as controls.

**Gene Transfer Vectors:** The recombinant adenovirus carrying the *Escherichia coli* β-galactosidase gene under the control of the human cytomegalovirus enhancer/promotor (Ad/CMV/β-gal) or the replication defective adenovirus were

propagated on 293 cells and purified and stored using techniques previously reported (Nguyen *et al.*, 1996). Viral titer was determined by UV spectrophotometric analysis (Nguyen *et al.*, 1996a; 1996b). To determine if the CDDP-induced enhancement of gene expression would be dependent on the method of gene delivery, two other gene delivery systems were used: the Ad/PLL/DNA complex (Nguyen *et al.*, 1996a; 1996b) and cationic liposomes (Lipofectamine, Gibco BRL). CDDP (0.062 µg/ml)-treated H1299 cells were incubated with either gene delivery system carrying a plasmid that contained the β-gal gene under the control of the CMV enhancer/promotor (CMV/β-gal) 48 hours after drug removal. Plasmid DNA was isolated by the alkaline lysis technique and purified by using Qiagen DNA purification kits (Qiagen, Chatsworth, CA). Plasmid containing liposome preparations and gene transfer were performed according to the manufacturer's protocol. The conjugated Ad/PLL/DNA complex was constructed and used to deliver the β-gal gene to H1299 and H460 cells as described elsewhere (Nguyen *et al.*, 1996a; 1996b).

**Quantitation of β-gal Gene Expression:** Adenoviral infection was performed 2 days after exposure to CDDP (0.062 µg/ml × 24 hours), with cell extracts being obtained 24 hours later and quantitatively assayed for the expression of the β-galactosidase gene using O-nitrophenyl β-d-galactoside (ONPG) as a substrate (MacGregor *et al.*, 1987). The magnitude of CDDP-induced enhancement of gene expression at a given MOI was determined by calculating the enhancement index:

$$\text{Enhancement index} = \frac{\beta\text{-gal activity of CDDP-treated cells}}{\beta\text{-gal activity of control cells}}$$

### ***In Vivo* CDDP-Induced Enhancement of β-gal Gene Expression**

Subcutaneous (SC) tumor nodules were created by injecting  $1 \times 10^7$  H1299 cells suspended in 100 µl of PBS into the dorsal flank SC tissue of nude mice (nu/nu, Charles River, Wilmington, MA) that had received 350 Rads of total body irradiation. Tumors, approximately 250 mm<sup>3</sup> in size, consistently formed within 3 to 4 weeks of tumor cell

implantation. Intratumoral injections of Ad/CMV/ $\beta$ -gal ( $5 \times 10^8$  viral particles) were done at 0, 2, 4, and 6 days following an intraperitoneal injection of CDDP (5  $\mu$ g/g body weight in 100  $\mu$ l of PBS) or PBS only (n = 4 per groups). The tumors were excised 48 hours after adenovirus injection, washed in cold PBS, embedded in tissue freezing medium (Fisher Scientific, Houston, TX) and snap-frozen by immersion in 2-methylbutane which was chilled over liquid nitrogen. Serial frozen sections (8  $\mu$ , thick) of the tumor mass obtained at 2-mm intervals were fixed and stained with X-gal as previously described (Ponder *et al.*, 1991) and then counterstained with nuclear fast red (PolyScientific, Bay Shore, NY). Quantitative assessment of  $\beta$ -gal gene expression (blue cells) was determined by digital image analysis (Samba 4000, Immuno Software, Version 3.0; Image Products International, Inc., Chantilly, VA) for each tumor section. *In vivo*  $\beta$ -gal gene transduction of tumor masses was calculated as the mean of the percentage of blue cells per surface area of representative sections. Tumor-bearing animals receiving intraperitoneal injection of PBS and subsequent intratumoral injection of Ad/CMV/ $\beta$ -gal at identical time points, served as controls. The magnitude of enhancement was calculated by dividing values from CDDP and Ad/CMV/ $\beta$ -gal treated tumors by values from Ad/CMV/ $\beta$ -gal control treated tumors.

### Statistical Analysis

The results are presented as means  $\pm$  standard deviations. Analysis of variance (ANOVA) and Student's *t*-test were used for statistical analysis;  $p < 0.05$  is considered significant.

## RESULTS

### Time Course and Dose Response Analysis

The initial response of incubating the H1299 and H460 cells with CDDP at 1 and 4  $\mu$ g/ml for 24 hours, ranged from a significant depression of growth to complete inhibition of proliferation; however, those cells were viable as demonstrated by trypan blue exclusion. At CDDP concentrations lower than 0.25  $\mu$ g/ml, cell growth remained

unchanged as compared to control cells populations. A brief 24-hour exposure of H1299 and H460 cells to CDDP (0.062  $\mu\text{g/ml}$ ) resulted in an increase of gene expression in target cells infected by Adv/CMV/ $\beta$ -gal as quantitated by X-gal staining (FIG. 1A). Maximal enhancement (2 to 2.5 fold increase in the percentage of positively stained cells as compared to concurrent controls) was noted when cells were infected with Adv/CMV/ $\beta$ -gal 48 hours after removal of CDDP ( $p < 0.001$ ). The enhancement effect was short-lived as the percentage of positively stained, treated cells, infected on day 4 or 5 after CDDP exposure was similar to that of the control. There was also a clear CDDP dose-related enhancement of  $\beta$ -gal gene expression in H1299 and H460 cells when infected with Adv/CMV/ $\beta$ -gal 48 hours after CDDP removal (FIG. 1B). Maximal enhancement occurred at a CDDP concentration of 0.062  $\mu\text{g/ml}$ , however, there was also an enhancement at 0.0039 and 1  $\mu\text{g/ml}$ . Optimal conditions for maximal enhancement of gene expression thus appears to be at the CDDP concentration of 0.062  $\mu\text{g/ml}$  and gene transfer at about 48 hours after drug treatment.

### Mechanism of Enhanced Gene Expression

To determine if prior incubation of target cells with CDDP would result in an increase in the ability of cells to take up gene-delivering vectors or an increase in the expression of successfully delivered genes,  $\beta$ -gal gene expression was quantitated in H1299 and H460 cell lines treated with CDDP and then infected with Ad/CMV/ $\beta$ -gal at increasing MOI's. If CDDP mediates an enhancement of adenovirus uptake as compared to non-treated controls, then there should be a proportional increase in enhancement indices with increasing MOI's. On the other hand, if prior CDDP exposure results in enhanced gene expression only, then the enhancement index should be independent of the increasing MOI. The analysis showed that the  $\beta$ -gal activity in H1299 infected cells increased as the MOI's were elevated from 5 to 100 (control H1299 cells of  $0.82 \pm 0.11$  to  $16.8 \pm 3.2$  U/mg protein and CDDP-treated H1299 cells of  $2.2 \pm 0.40$  to  $42.0 \pm 2.6$  U/mg protein), indicating unchanged enhancement indices (3 to 2.5) with increasing MOI's (FIG. 2A). Similarly,  $\beta$ -gal gene expression in CDDP-treated H460 cells increased from  $0.28 \pm 0.06$  to  $12.2 \pm 2.1$  U/mg protein, as compared to  $\beta$ -gal gene expression in control H460 cells of  $0.056 \pm 0.016$  to  $7.6 \pm 1.8$  U/mg protein. The enhancement indices in H460

cells, therefore, decreased from a value of 5 at an MOI of 5, to a value of 1.6 at an MOI of 100 (FIG. 2B). This observation implies that CDDP induces an increase in the expression of successfully delivered genes and not an increase in adenovirus uptake.

## 5    **Enhanced $\beta$ -gal Gene Expression by DNA-Damaging Agents**

Infection of H1299 cells that were similarly treated with other antineoplastic agents such as vincristine, methotrexate, 5-fluorouracil or transplatinum with Ad/CMV/ $\beta$ -gal did not result in enhanced gene expression over a wide range of drug concentrations (FIG. 3).  
10    Methotrexate and 5-FU are anti-metabolites which interfere with synthesis of nucleic acid precursors; vincristine binds to and inhibits microtubular formation and impairs cellular mitosis (Fritsch *et al.*, 1993). On the other hand, exposure of H1299 cells to the DNA-damaging agents VP-16 (etoposide) and ionizing irradiation (at 4 and 8 Grey) enhanced gene expression in tumor cells infected with Adv/CMV/ $\beta$ -gal (FIG. 3). These  
15    pharmacological agents induce DNA strand breaks by different mechanisms: CDDP causes intra- and inter-strand crosslinking of DNA by forming guanine adducts and eventually DNA strand breaks (Eastman, 1990); Etoposide forms a tertiary complex between DNA and topoisomerase II and eventually produces protein-linked DNA single and double strand breaks (Kaufman, 1989); ionizing irradiation stimulates production of  
20    oxygen free radicals which react with macromolecules and induces DNA damage (Cole *et al.*, 1980).

## **Enhanced $\beta$ -gal Gene Expression in Other Cell Types and With Different Vectors**

25    When CDDP-treated H1299 cells were incubated with 2 other gene delivery systems (the conjugated Ad/PLL/DNA complex or the cationic phospholipid lipofectamine),  $\beta$ -gal gene expressing cells occurred at the same CDDP doses and to the same magnitude as observed following adenovirus-mediated gene transfer (FIG. 4A). This indicates that the enhancement by CDDP is not vector dependent. This CDDP-induced  
30    enhancement of transgene expression following Ad/CMV/ $\beta$ -gal infection was also observed in other malignant cell lines tested under the same experimental conditions (FIG. 4B). The magnitude of maximal enhancement was dependent on the cell line tested

(ranging from 1.6-fold (A549) to 2.7-fold (H1299)), but it consistently occurred when gene transfer was performed 48 hours after exposure to 0.062  $\mu\text{g/ml}$  of CDDP. Similar exposure of stationary primary human bronchial epithelial cells (90% confluent in culture) to a low concentration (0.062  $\mu\text{g/ml}$ ) of CDDP failed to enhance gene expression in these  
5 normal cells after Ad/CMV/ $\beta$ -gal infection (FIG. 4B). However, at higher drug concentrations (1 and 4  $\mu\text{g/ml}$ ), there was a dose-dependent elevation of the percentage of  $\beta$ -gal-positive cells that was 1.5- to 2.5-fold higher than that observed in unexposed NHBE cells.

## 10 *In Vivo* Enhancement of $\beta$ -gal Gene Expression

To analyze the effect of CDDP on enhancing gene expression in tumor cells *in vivo*, a time course analysis was done. The time course of enhanced  $\beta$ -gal expression in H1299 cells *in vivo* was similar to the *in vitro* time course (FIG. 5). The overall gene  
15 expression in subcutaneous H1299 tumors of animals injected with PBS and Ad/CMV/ $\beta$ -gal slightly decreased ( $29.2 \pm 9.0\%$  on day 0 to  $18.2 \pm 5.6\%$  on day 6) over time as tumors continued to grow while the viral titer remained unchanged. There was no change in the  $\beta$ -gal gene expression in H1299 cells when CDDP and Ad/CMV/ $\beta$ -gal were concurrently administered (day 0). However, the percentages of positively stained cells in  
20 tumors injected with Ad/CMV/ $\beta$ -gal 2 and 4 days after CDDP administration were 1.6- to 2.8-fold higher than the values of the respective controls ( $46.0 \pm 4.3$  vs  $28.7 \pm 6.2$  on day 2 and  $70.2 \pm 11$  vs.  $25.0 \pm 7.0$  on day 4,  $p < 0.01$ ). When gene transfer was performed 6 days after CDDP administration, the increase in expression was slightly higher than that of the control tumors ( $23.4 \pm 5$  vs.  $18.2 \pm 5.6$ ,  $p$  (FIG. 5).

25

**EXAMPLE 2: GENE THERAPY STRATEGY FOR HUMAN NON-SMALL CELL  
LUNG CANCER: COMBINATION OF SEQUENTIAL CISPLATIN  
ADMINISTRATION AND ADENOVIRUS-MEDIATED p53 GENE TRANSFER**

**MATERIALS AND METHODS**

**Cells and Culture Conditions**

The human NSCLC cell line H1299 with a homozygous deletion of p53 gene was grown in RPMI-1640 medium supplemented with 5% heat-inactivated fetal calf serum (FCS), 10 mM glutamine, 100 units/ml of penicillin, 100 µg/ml of streptomycin, 0.25 µg/ml of amphotericin B (Gibco-BRL, Grand Island, NY) (RPMI-complete). The H322 cell line was maintained in RPMI-complete medium supplemented with 10% FCS. The 293 cells were maintained in complete high glucose Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat-inactivated FCS.

**Recombinant Adenovirus Production**

The properties of the adenovirus Ad/CMV/p53 have previously been demonstrated in the inventors' laboratory (Diller *et al.*, 1990). The replication-defective E1A-deleted adenovirus d1312 was used as a control. The virus was amplified on a large scale on 293 cells according to methods previously reported (Zhang *et al.*, 1995a). Viral titer was determined by UV-spectrophotometric analysis (viral particles/ml). The absence of replicative competence adenovirus was verified by a PCR™ technique previously reported by Zhang *et al.* (1995b). The purified virus was stored in 10% glycerol at -80°C. Infection of H1299 cells was carried out by dilution of the viral stock to appropriate concentrations, followed by the addition of viral solutions to cell monolayers incubated in 1 ml of RPMI-1640 medium containing 2% FCS. The cells were incubated for 2 hrs at 37°C in a 5% CO<sub>2</sub> incubator, after which 2 ml of RPMI-1640 complete medium supplemented with 5% FCS was added into the wells.

### Cell Proliferation Assay

H1299 cells were exposed to CDDP (0.062  $\mu$ g/ml of medium) for 24 hrs. Cells were then washed twice with PBS, trypsinized and seeded in 6-well plates ( $10^5$  cells/well).  
5 Forty eight hours later, cells were infected with Ad/CMV/p53 at the multiplicity of infection (MOI) of 5. Daily cell counts were performed for 5 days following transfection to study the tumor suppression effect of the combination of sequential CDDP and adenovirus-mediated p53 gene transfer. Controls consisted of untreated cells or cells exposed to CDDP only, cells transfected with Ad/CMV/p53 only and cells transfected  
10 with d1312 (similar MOI) with or without prior CDDP treatment.

### Western Blot Analysis

H1299 cells were infected with Ad/CMV/p53 (MOI = 1) 48 hrs after CDDP  
15 exposure. Cells were then harvested at 6, 12, 24, 36 and 48 hrs after transfection for Western blot analysis of p53 protein expression, as described elsewhere (Diller *et al.*, 1990). Unexposed cells similarly transfected with Ad/CMV/p53 were used as controls. The human NSCLC cell line H322 which overexpresses mutant p53 protein was used as a positive control. Relative quantities of exogenous p53 protein were determined by  
20 densitometer (Molecular Dynamics, Sunnyvale, CA).

### p53 Immunocytochemical Staining

The infected cells were fixed 3.8% formalin at 12 and 24 hrs after transfection with  
25 Adv/CMV/p53 (MOI = 5) with or without prior CDDP exposure and treated with 3%  $H_2O_2$  in methanol for 5 minutes. The Vectastain kit (Vector Laboratories, Burlingana, CA) was used for the immunocytochemical staining. The primary antibody used was the mouse monoclonal anti-p53 antibody Pab 1801 (Oncogene Science, Manhasset, NY). The secondary antibody was the biotinylated anti-mouse IgG antibody (Vector Laboratories).  
30 The avidin-biotin-peroxidase complex (ABC) kit was used to detect the antigen-antibody complex. The cells were counterstained with Harris Hematoxylin (Sigma, St. Louis, MO) and mounted with cover slips using Cytoseal 60.



### ***In Situ* TUNEL Assay for Apoptosis**

CDDP-treated or control H1299 cells were fixed in 50% acetone/ethanol for 20  
5 minutes at -20°C 12 hrs and 24 hrs after transfection with Adv/CMV/p53 (MOI = 5). *In situ* TdT (terminal deoxynucleotidyl transferase)-mediated dUTP-biotin nick end-labeling (TUNEL) assay was performed according to the procedure described elsewhere (Gavrieli *et al.*, 1992). H1299 cells used as positive controls were treated with DNase I (Gibco BRL, Gaithersburg, MD), for 1 hr at 37°C (50 µg/ml in 10 mM Tris-HCl pH = 7.5; 140  
10 mM sodium cacodylate, 4 mM MgCl<sub>2</sub>, 10 mM dithiothreitol).

### ***In Vivo* Combination of Sequential CDDP and Adv/CMV/p53 Administrations**

H1299 tumor xenografts were created by injecting  $1 \times 10^7$  cells suspended in 100  
15 µl of PBS into the dorsal flank subcutaneous space of nude mice that had received 350 Rads of total body irradiation prior to injection. Subcutaneous tumor nodules of 200 to 250 mm<sup>3</sup> in size were formed 3 to 4 weeks later. Different strategies of CDDP and Ad/CMV/p53 combination were studied for their tumoricidal efficacy: a) intraperitoneal (ip) CDDP (5 µg/g body weight) given on day 0 and followed by intratumoral  
20 Ad/CMV/p53 injections of  $1.5 \times 10^{10}$  viral particles/100 µl PBS in a single dose on day 3 or in 3 equally divided doses of  $5 \times 10^9$  viral particles/100 µl PBS on days 2, 4 and 6; b) simultaneous CDDP and Ad/CMV/p53 administrations in single or 3 equally divided doses (divided CDDP doses: 1.67 µg/g body weight) and c) ip CDDP given 3 days after  
25 completion of intratumoral Ad/CMV/p53 injections. The divided dose regime was designed to address issues that may limit the use of high virus titer and volume of injectate such as toxicity and the low titer of viral stock. The controls consisted of tumors injected with either PBS, Ad/CMV/p53 without prior systemic CDDP, d1312 with or without prior  
30 ip CDDP or tumor-bearing animals receiving ip CDDP only. Tumor sizes were measured every 2 days for 32 days and tumor volumes were estimated by assuming a spherical shape with the average tumor diameter calculated as the square root of the product of the orthogonal diameters. All animals were treated according to guidelines developed by the

M.D. Anderson Animal Care and Use Committee. All mice were sacrificed when tumors grew to 4000 mm<sup>3</sup>.

### Statistical Analysis

Results were presented as mean  $\pm$  standard deviation. Analysis of variance (ANOVA) and two-tailed Student's *t* test were used for statistical analysis of multiple groups and pair-wise comparison respectively,  $p < 0.05$  is considered significant.

## RESULTS

### *In Vitro* Proliferation Assay

H1299 cells were treated with a combination of sequential CDDP and Ad/CMV/p53 infection and then analyzed by cell proliferation assay to determine if the combination of CDDP and adenovirus-mediated p53 gene transfer would result in a superior growth-inhibiting effect. Treating H1299 cells with CDDP (0.0625  $\mu$ g/ml) for 24 hrs had no effect on cell growth *in vitro*, nor did mock infection with d1312 with or without prior CDDP treatment. On the other hand, exposure of H1299 cells to CDDP 48 hrs prior to Adv/CMV/p53 transfection resulted in 61% and 55% increased inhibition of tumor proliferation on day 3 and 5 after p53 transfer when compared to control H1299 cells similarly transfected with Adv/CMV/p53 (FIG. 6).

### Western Blot Analysis and p53 Immunocytochemical Staining

The effect of CDDP treatment on levels of p53 protein was determined in CDDP-treated and untreated control cells by Western blotting of cell lysates harvested at different time points after Adv/CMV/p53 infection. Expression of the p53 gene occurred as early as 6 hours after Adv/CMV/p53 transfection in both CDDP-treated and control H299 cells. However, prior exposure to CDDP led to a higher level of p53 at 12, 24, 36 and 48 hrs after transfection. Densitometry analysis showed that the relative levels of p53 protein (normalized for  $\beta$ -actin levels) in CDDP-treated cells were 0.73, 1.39, 1.49, 1.20

compared to 0.23, 0.90, 0.68, 0.62 (for a 2- to 3-fold increase in the levels of p53 protein) in cells without prior CDDP at each of the time points studied. The p53 levels at 6 hrs after transfection were too low to be analyzed. CDDP treatment appeared to significantly increase the p53 gene expression but did not seem to alter the transduction kinetics of H1299 cells. Immunocytochemical staining of H1299 cells 24 hrs after p53 gene transfer demonstrated that almost all of the CDDP-treated cells stained positive, at a significantly higher intensity, for the p53 protein as compared to cells not exposed to CDDP. This elevation of p53 gene expression correlates well with the enhanced tumoricidal effect of the combination of CDDP and Adv/CMV/p53 *in vitro* as demonstrated in FIG. 6.

### Induction of Apoptosis Following Adv/CMV/p53 Transfection

The *in situ* TUNEL assay was used to analyze the extent of apoptosis that occurred in Ad/CMV/p53-transfected H1299 cells. In cells that were not exposed to CDDP prior to gene transfer, apoptosis began 12 hrs after p53 transfection with few TUNEL-positive cells being visualized with significant apoptosis was detected 24 hrs after Ad/CMV/p53 transfection. Apoptotic cells could be readily seen in CDDP-treated cells as early as 12 hrs after transfection, at a much higher frequency than in untreated cells. By 24 hrs, almost all CDDP-treated cells stained positive for DNA fragmentation. The degree of apoptosis and the timing of its occurrence follow very closely the time course of p53 gene expression in CDDP-treated cells. Thus enhanced and accelerated p53 gene expression in cells treated with sequential CDDP and Ad/CMV/p53 most probably resulted in an early and intense induction of apoptosis which translated to an increased tumoricidal effect.

### Inhibition of Tumor Growth *In Vivo*

**Divided-Dose Regime:** Tumors grew rapidly in groups of control animals receiving only ip CDDP or intratumoral PBS with tumor sizes reaching the maximal allowable volume of 4000 mm<sup>3</sup> 22 to 26 days after treatment. Tumors injected with d1312 with or without prior ip CDDP showed some degree of growth retardation (up to 30% of normal tumor growth) secondary to vector toxicity. Injections of Ad/CMV/p53 ( $1.5 \times 10^{10}$  viral particles in 3 equally divided doses) resulted in inhibition of tumor growth during and

immediately after the treatment. These tumors, however, resumed growth at the normal rate, 10 days after the last injection; reaching the mean tumor volume of  $3357 \pm 391 \text{ mm}^3$  32 days after the onset of therapy. Intraperitoneal CDDP administration given 2 days prior to the beginning of the gene therapy schedule resulted in a more pronounced inhibition of tumor development. There was a regression of tumor mass that lasted for 14 days before the tumor growth resumed which was at a much slower rate. At the end of the observation period, the average tumor size was  $1497 \pm 221 \text{ mm}^3$  ( $p < 0.001$  vs Ad/CMV/p53 without prior CDDP). Systemic CDDP administration prior to p53 gene replacement therapy, therefore, resulted in a synergistic effect of tumor growth inhibition that was responsible for at least a 55.4% further reduction in tumor size.

**Single-Dose Regime:** When Ad/CMV/p53 was given in a single dose, inhibition of tumor growth was greater than that seen with the divided-dose regime:  $1936 \pm 308$  vs  $3357 \pm 391$  (without CDDP) and  $773 \pm 197$  vs  $1497 \pm 221$  (with prior CDDP) respectively. The combination of CDDP administration 3 days prior to a single intratumoral injection of Ad/CMV/p53 ( $1.5 \times 10^{10}$  viral particles) resulted in a 60% further suppression of tumor growth compared to tumors treated with Ad/CMV/p53 without prior CDDP. To illustrate the importance of the timing of CDDP administration in relation to p53 gene transfer, CDDP ( $5 \mu\text{g/ml}$ ) was given at the same time as Ad/CMV/p53 ( $1.5 \times 10^{10}$  viral particles) injections in a second group of mice, and in a third group CDDP was given 3 days after adenovirus-mediated p53 gene transfer. The combination of sequential CDDP and Ad/CMV/p53 showed the most significant tumor growth inhibition effect with the mean tumor volume of  $773 \pm 197 \text{ mm}^3$ , compared to  $1257 \pm 225 \text{ mm}^3$ , when CDDP was given concurrently with adenovirus injections, and  $1750 \pm 214 \text{ mm}^3$ , when CDDP was given after adenovirus-mediated p53 transfer ( $p < 0.001$  by ANOVA). Ad/CMV/p53 significantly prolonged the survival of treated animals to  $62.8 \pm 4.6$  days (range: 60 to 70 days) after the onset of therapy compared to tumor-bearing mice injected with Ad/CMV/p53 alone of  $46.0 \pm 4.4$  days or untreated controls of  $23.4 \pm 2.9$  days ( $p < 0.001$ ).

H. References

The following references, to the extent that they provide exemplary procedural details or other information supplementary to that set forth herein, are incorporated by reference.

Anderson *et al.*, U.S. Patent 5,399,346.

Baichwal and Sugden, "Vectors for gene transfer derived from animal DNA viruses: Transient and stable expression of transferred genes," *In*: Kucherlapati R, ed. Gene transfer. New York: Plenum Press, pp. 117-148, 1986.

Bargonetti *et al.*, *Cell*, 65:1083-1091, 1991.

Benvenisty and Neshif, "Direction introduction of genes into rats and expression of the genes," *Proc. Nat'l Acad. Sci. USA*, 83:9551-9555, 1986.

Bittner *et al.*, *Methods in Enzymol.*, 153:516-544, 1987.

Casey *et al.*, "Growth suppression of human breast cancer cells by the introduction of a wild-type p53 gene," *Oncogene*, 6:1791-1797, 1991.

Chang *et al.*, "Foreign gene delivery and expreddion in hepatocytes using a hepatitis B virus vector," *Hepatology*, 14:134A, 1991.

Chen and Okayama, "High-efficiency transfection of mammalian cells by plasmid DNA," *Mol. Cell Biol.*, 7:2745-2752, 1987.

Coffin, "Retroviridae and their replication," *In*: *Virology*, Fields *et al.* (eds.), New York: Raven Press, pp. 1437-1500, 1990.

Colberre-Garapin *et al.*, 1981.

- Cole *et al.*, In: *Mechanisms of Injury*, eds. Meyn, R.E. & Withers, H.R., 1980.
- 5 Couch *et al.*, "Immunization with types 4 and 7 adenovirus by selective infection of the  
intestinal tract," *Am. Rev. Resp. Dis.*, 88:394-403, 1963.
- Coupar *et al.*, "A general method for the construction of recombinant vaccinia virus  
expressing multiple foreign genes," *Gene*, 68:1-10, 1988.
- 10 DeVita *et al.*, "Cancer Principles and Practice of Oncology," 3rd. ed.
- Diller *et al.*, "p53 Functions as a Cell Cycle Control Protein in Osteosarcomas," *Mol. Cell  
Biol.*, 10(11):5772-5781, 1990.
- 15 Dubensky *et al.*, "Direct transfection of viral and plasmid DNA into the liver or spleen of  
mice," *Proc. Nat. Acad. Sci. USA*, 81:7529-7533, 1984.
- Eastman, *Cancer Cell*, 2:275-280, 1990.
- 20 Fechheimer *et al.*, "Transfection of mammalian cells with plasmid DNA by scrape loading  
and sonication loading," *Proc. Nat'l Acad. Sci. USA*, 84:8463-8467, 1987.
- Ferkol *et al.*, Ferkol *et al.*, *FASEB J.*, 7:1081-1091, 1993.
- 25 Fields *et al.*, *Science*, 249:1046-1049, 1990.
- Fraley *et al.*, "Entrapment of a bacterial plasmid in phospholipid vesicles: Potential for gene  
transfer," *Proc. Nat'l Acad. Sci. USA*, 76:3348-3352, 1979.
- 30 Freshner, "Animal Cell Culture: A Practical Approach," Second Edition, Oxford/New York,  
IRL Press, Oxford University Press, 1992.

Friedmann, "Progress toward human gene therapy," *Science*, 244:1275-1281, 1989.

Fritsch *et al.*, *Oncogene*, 8:307-318, 1993.

- 5 Gavrieli *et al.*, "Identification of Programmed Cell Death *In Situ Via* Specific Labeling of Nuclear DNA Fragmentation," *J. Cell Biol.*, 119:493-501, 1992.

Ghosh-Choudhury, *et al.*, "protein IX, a minor component of the human adenovirus capsid, is essential for the packaging of full length genomes," *EMBO J.*, 1733-1739, 1987.

10

Ghosh and Bachhawat, "Targeting of liposomes to hepatocytes," *In: Wu G. and C. Wu ed. Liver diseases, targeted diagnosis and therapy using specific receptors and ligands.* New York: Marcel Dekker, pp. 87-104, 1991.

- 15 Gomez-Foix *et al.*, "Adenovirus-mediated transfer of the muscle glycogen phosphorylase gene into hepatocytes confers altered regulation of glycogen," *J. Biol. Chem.*, 267:25129-25134, 1992.

20 Goodman and Gilman's "The Pharmacological Basis of Therapeutics." 8th ed. Pergammon Press, New York, 1990.

Gopal, "Gene transfer method for transient gene expression, stable transfection, and cotransfection of suspension cell cultures," *Mol. Cell Biol.*, 5:1188-1190, 1985.

- 25 Graham, *et al.*, "Characteristics of a human cell line transformed by DNA from human adenovirus type 5," *J. Gen Virol.* 36:59-72, 1977.

30 Graham and Prevec, "Manipulation of adenovirus vectors." *In: E.J. Murray (ed.), Methods in Molecular Biology, Gene Transfer and Expression Protocols*, New Jersey: The Humana Press Inc., 109-128, 1991.

Graham and Van Der Eb, "A new technique for the assay of infectivity of human adenovirus 5 DNA", *Virology*, 52:456-467, 1973.

5 Grunhaus and Horwitz, "Adenoviruses as cloning vectors," *Semin. Virology*, 3:237-2542, 1992.

Harland and Weintraub, "Translation of mammalian mRNA injected into *Xenopus* oocytes is specifically inhibited by antisense RNA," *J. Cell Biol.*, 101:1094-1099, 1985.

10 Hermonat and Muzycska, "Use of adenoassociated virus as a mammalian DNA cloning vector: Transduction of neomycin resistance into mammalian tissue culture cells," *Proc. Nat'l Acad. Sci. USA*, 81:6466-6470, 1984.

15 Hersdorffer *et al.*, "Efficient gene transfer in live mice using a unique retroviral packaging line," *DNA Cell Biol.*, 9:713-723, 1990.

Herz and Gerard, "Adenovirus-mediated transfer of low density lipoprotein receptor gene acutely accelerates cholesterol clearance in normal mice," *Proc. Nat'l Acad. Sci. USA* 90:2812-2816, 1993.

20 Hollstein *et al.*, "p53 mutations in human cancers," *Science*, 253:49-53, 1991.

Horwich *et al.*, "Synthesis of hepadenovirus particles that contain replication-defective duck hepatitis B virus genomes in cultured HuH7 cells," *J. Virol.*, 64:642-650, 1990.

25 Jones and Shenk, "Isolation of deletion and substitution mutants of adenovirus type 5," *Cell*, 13:181-188, 1978.

30 Kaneda *et al.*, "Increased expression of DNA cointroduced with nuclear protein in adult rat liver," *Science*, 243:375-378, 1989.

Karlsson *et al.*, *EMBO J.*, 5:2377-2385, 1986.



Kato *et al.*, "Expression of hepatitis  $\beta$  virus surface antigen in adult rat liver," *J. Biol. Chem.*, 266:3361-3364, 1991.

5 Kaufman, *Cancer Res.*, 49:5870-5878, 1989.

Klein *et al.*, "High-velocity microprojectiles for delivering nucleic acids into living cells," *Nature*, 327:70-73, 1987.

10 Le Gal La Salle *et al.*, "An adenovirus vector for gene transfer into neurons and glia in the brain," *Science* 259:988-990, 1993.

Levrero *et al.*, "Defective and nondefective adenovirus vectors for expressing foreign genes *in vitro* and *in vivo*," *Gene*, 101:195-202, 1991.

15

MacGregor *et al.*, *Somat Cell Mol. Genet.*, 13:253-265, 1987.

Mann *et al.*, "Construction of a retrovirus packaging mutant and its use to produce helper-free defective retrovirus," *Cell*, 33:153-159, 1983.

20

Markowitz *et al.*, "A safe packaging line for gene transfer: Separating viral genes on two different plasmids," *J. Virol.*, 62:1120-1124, 1988.

Mercer, "Cell cycle regulation and the p53 tumor suppressor protein," *Critic. Rev. Eukar.*

25

*Gene Express.* 2:251-263, 1992.

Mietz *et al.*, *EMBO*, 11:5013,5020, 1992.

Mizrahi, "Production of human interferons: An overview," *Process Biochem.*, (August):9-12, 1983.

30

Mulligan, "The Basic Science of Gene Therapy," *Science*, 260:926-932, 1993.

Nguyen *et al.*, *J. Virol.*, (submitted), 1996a.

Nguyen *et al.*, *Cancer Gene Ther.*, (submitted), 1996b.

5

Nicolas and Rubenstein, "Retroviral vectors," *In: Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt (eds.), Stoneham: Butterworth, pp. 494-513, 1988.

10 Nicolau and Sene, "Liposome-mediated DNA transfer in eukaryotic cells," *Biochem. Biophys. Acta*, 721:185-190, 1982.

Nicolau *et al.*, "Liposomes as carriers for *in vivo* gene transfer and expression," *Methods Enzymol.*, 149:157-176, 1987.

15

Paskind *et al.*, "Dependence of moloney murine leukemia virus production on cell growth," *Virology*, 67:242-248, 1975.

Perales *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:4086-4090, 1994.

20

Phillips *et al.*, *In: Large Scale Mammalian Cell Culture* (Feder, J. and Tolbert, W. R., eds.), Academic Press, Orlando, FL, U.S.A., 1985.

Ponder *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:1217-1221, 1991.

25

Potter *et al.*, "Enhancer-dependent expression of human k immunoglobulin genes introduced into mouse pre-B lymphocytes by electroporation," *Proc. Nat'l Acad. Sci. USA*, 81:7161-7165, 1984.

30 Ragot *et al.*, "Efficient adenovirus-mediated transfer of a human minidystrophin gene to skeletal muscle of mdx mice," *Nature*, 361:647-650, 1993.

Renan, "Cancer genes: current status, future prospects, and applicants in radiotherapy/oncology," *Radiother. Oncol.*, 19:197-218, 1990.

Rich *et al.*, "Development and analysis of recombinant adenoviruses for gene therapy of cystic fibrosis," *Hum. Gene Ther.*, 4:461-476, 1993.

Ridgeway, "Mammalian expression vectors," *In: Rodriguez RL, Denhardt DT, ed. Vectors: A survey of molecular cloning vectors and their uses. Stoneham: Butterworth, pp. 467-492, 1988.*

Rippe *et al.*, "DNA-mediated gene transfer into adult rat hepatocytes in primary culture," *Mol. Cell Biol.*, 10:689-695, 1990.

Rosenfeld *et al.*, "Adenovirus-mediated transfer of a recombinant  $\alpha$ 1-antitrypsin gene to the lung epithelium *in vivo*," *Science*, 252:431-434, 1991.

Rosenfeld *et al.*, "*In vivo* transfer of the human cystic fibrosis transmembrane conductance regulator gene to the airway epithelium," *Cell*, 68:143-155, 1992.

Roux *et al.*, "A versatile and potentially general approach to the targeting of specific cell types by retroviruses: Application to the infection of human cells by means of major histocompatibility complex class I and class II antigens by mouse ecotropic murine leukemia virus-derived viruses," *Proc. Natl. Acad. Sci. USA*, 86:9079-9083, 1989.

Russell *et al.*, *Proc. Natl. Acad. Sci. USA*, 92:5719-5723, 1995.

Shaw *et al.*, 89:4495-4499, 1992.

Son & Huang, *Proc. Natl. Acad. Sci. USA*, 91:12669-12672, 1994.

- Stratford-Perricaudet and Perricaudet, "Gene transfer into animals: the promise of adenovirus," p. 51-61, *In: Human Gene Transfer*, Eds, O. Cohen-Haguenauer and M. Boiron, Editions John Libbey Eurotext, France, 1991.
- 5 Stratford-Perricaudet *et al.*, "Evaluation of the transfer and expression in mice of an enzyme-encoding gene using a human adenovirus vector," *Hum. Gene Ther.*, 1:241-256, 1990.
- 10 Takahasi *et al.*, "Wild-type but not mutant p53 suppresses the growth of human lung cancer cells bearing multiple genetic lesions," *Cancer Res.*, 52:2340-2342, 1992.
- Temin, "Retrovirus vectors for gene transfer: Efficient integration into and expression of exogenous DNA in vertebrate cell genome," *In: Gene Transfer*, Kucherlapati (ed.), New York: Plenum Press, pp. 149-188, 1986.
- 15 Top *et al.*, "Immunization with live types 7 and 4 adenovirus vaccines. II. Antibody response and protective effect against acute respiratory disease due to adenovirus type 7," *J. Infect. Dis.*, 124:155-160, 1971.
- 20 Tur-Kaspa *et al.*, "Use of electroporation to introduce biologically active foreign genes into primary rat hepatocytes," *Mol. Cell Biol.*, 6:716-718, 1986.
- Wagner *et al.*, *Science*, 260:1510-1513, 1990.
- 25 Weinberg, "Tumor suppressor gene," *Science*, 254:1138-1145, 1991.
- Wilcock and Lane, *Nature*, 349:429-431, 1991.
- Wong *et al.*, "Appearance of  $\beta$ -lactamase activity in animal cells upon liposome mediated gene transfer," *Gene*, 10:87-94, 1980.
- 30

Wu and Wu, "Receptor mediated *in vitro* gene transfections by a soluble DNA courier system," *J. Biol. Chem.*, 262:4429-4432, 1987.

5 Wu and Wu, "Evidence for targeted gene delivery to HepG2 hepatoma cells *in vitro*," *Biochemistry*, 27:887-892, 1988.

Wu and Wu, *Adv. Drug Delivery Rev.*, 12:159-167, 1993.

10 Yang *et al.*, "In vivo and *in vitro* gene transfer to mammalian somatic cells by particle bombardment," *Proc. Nat'l Acad. Sci. USA*, 87:9568-9572, 1990.

Yonish-Rouach *et al.*, *Nature*, 352:345-347, 1991.

15 Varmus *et al.*, "Retroviruses as mutagens: Insertion and excision of a non-transforming provirus alter the expression of a resident transforming provirus," *Cell*, 25:23-36, 1981.

Zakut-Houri *et al.*, *EMBO J.*, 4:1251-2355, 1985.

20 Zelenin *et al.*, "High-velocity mechanical DNA transfer of the chloramphenicol acetyltransferase gene into rodent liver, kidney and mammary gland cells in organ explants and *in vivo*," *FEBS Lett.*, 280:94-96, 1991.

25 Zhang *et al.*, "Detection of Wild-type Contamination in a Recombinant Adenoviral Preparation by PCR™," *BioTechniques*, 18:5-8, 1995b.

Zhang *et al.*, "Safety Evaluation of Ad5CMV-p53 *In Vitro* and *In Vivo*," *Human Gene Ther.*, 6:155-164, 1995a.

30

CLAIMS

1. A method for enhancing the expression of a transgene comprising:

5 (a) contacting a target cell with a DNA-damaging agent;

(b) removing said DNA-damaging agent from said target cell; and

10 (c) transferring said transgene into said target cell between about 1-3 days after removing said DNA-damaging agent.

2. The method of claim 1, wherein said target cell is a dividing cell.

15 3. The method of claim 2, wherein said target cell is a tumor cell.

4. The method of claim 3, wherein said tumor cell is cisplatin sensitive.

20 5. The method of claim 3, wherein said tumor cell is cisplatin insensitive.

25 6. The method of claim 1, wherein said DNA-damaging agent is selected from the group consisting of cisplatin, carboplatin, VP16, teniposide, daunorubicin, doxorubicin, dactinomycin, mitomycin, plicamycin, bleomycin, procarbazine, nitrosourea, cyclophosphamide, bisulfan, melphalan, chlorambucil, ifosfamide, merchlorhetamine, and ionizing radiation.

30

7. The method of claim 1, wherein said transgene is transferred at about 2 days after removing said DNA-damaging agent.

5           8. The method of claim 1, wherein said transfer of said transgene is accomplished by a technique selected from the group consisting of liposome-mediated transfection, receptor-mediated internalization and viral infection.

10           9. The method of claim 1, wherein said transgene is a tumor suppressor.

10. The method of claim 9, wherein said tumor suppressor is p53.

15           11. The method of claim 10, wherein said p53 transgene is under the transcriptional control of a promoter.

20           12. The method of claim 11, wherein said promoter is the CMV IE promoter.

25           13. The method of claim 12, wherein said transgene is regulated by a polyadenylation signal.

30           14. The method of claim 13, wherein said polyadenylation signal is an SV40 polyadenylation signal.

15. The method of claim 14, wherein said p53 transgene is carried in an adenoviral vector.

1/11




FIG. 1A

SUBSTITUTE SHEET (RULE 26)



2/11




FIG. 1B

SUBSTITUTE SHEET (RULE 26)

3/11




FIG. 2A

SUBSTITUTE SHEET (RULE 26)

4/11




FIG. 2B

SUBSTITUTE SHEET (RULE 26)

5/11




FIG. 3

SUBSTITUTE SHEET (RULE 26)

6/11





FIG. 4A

SUBSTITUTE SHEET (RULE 26)

7/11



CELL LINES

FIG. 4B

SUBSTITUTE SHEET (RULE 26)

8/11




FIG. 5

SUBSTITUTE SHEET (RULE 26)

9/11




FIG. 6

SUBSTITUTE SHEET (RULE 26)



10/11




FIG. 7A

SUBSTITUTE SHEET (RULE 26)

11/11




FIG. 7B

SUBSTITUTE SHEET (RULE 26)

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 97/05325

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/87 C12N15/86 C12N15/67 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PROCEEDINGS OF THE AMERICAN ASSOCIATION OF CANCER RESEARCH, vol. 37, March 1996, pages 347-348, XP002039691 NGUYEN,D., ET AL. : "GENE THERAPY FOR LUNG CANCER: ENHANCEMENT OF TUMOR SUPPRESSION BY A COMBINATION OF SYSTEMIC CISPLATIN AND ADENOVIRUS-MEDIATED p53 GENE TRANSFER" see the whole document --- -/--	1-4,6-15

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 September 1997

Date of mailing of the international search report

16.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Holtorf, S

# INTERNATIONAL SEARCH REPORT

Int. Application No  
PCT/US 97/05325

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PROCEEDINGS OF THE AMERICAN ASSOCIATION OF CANCER RESEARCH, vol. 37, March 1996, page 347 XP002039692 NGUYEN, D., ET AL. : "ENHANCEMENT OF GENE TRANSDUCTION IN HUMAN CARCINOMA CELLS BY DNA-DAMAGING AGENTS" see the whole document	1-4,6-8
A	--- PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 91, December 1994, pages 12669-12672, XP002039693 SON, K. AND HUANG, L.: "EXPOSURE OF THE HUMAN OVARIAN CARCINOMA TO CISPLATIN TRANSIENTLY SENSITIZES THE TUMOR CELLS FOR LIPOSOME-MEDIATED GENE TRANSFER" cited in the application see the whole document	1-15
A	--- WO 95 33824 A (HUTCHINSON FRED CANCER RES ;ALEXANDER IAN E (US); RUSSELL DAVID W) 14 December 1995 page 2, line 35 - page 4, line 10; page 5, line 35; example 1	1-15
A	--- PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 92, June 1995, pages 5719-5723, XP002039694 RUSSELL, D.W., ET AL.: "DNA SYNTHESIS AND TOPOISOMERASE INHIBITORS INCREASE TRANSDUCTION BY ADENO-ASSOCIATED VECTORS" cited in the application see the whole document	1-15
P,X	--- THORACIC AND CARDIOVASCULAR SURGERY, vol. 112, no. 5, November 1996, pages 1372-1377, XP002039695 NGUYEN, D.M., ET AL. : "GENE THERAPY FOR LUNG CANCER: ENHANCEMENT OF TUMOR SUPPRESSION BY A COMBINATION OF SEQUENTIAL SYSTEMIC CISPLATIN AND ADENOVIRUS-MEDIATED p53 GENE TRANSFER" see the whole document -----	1-4, 6-11,13, 15

information on patent family members

PCr/US 97/05325

Form PCT/ISA/210 (patent family annex) (July 1992)

**THIS PAGE BLANK (USPTO)**